

University of Groningen

Synthesis of Carbapenems with Carbon Substituents at C-3 Using a Wittig Approach

Vries, Johannes G. de; Hauser, Gerhard; Sigmund, Gerhard

Published in:
Tetrahedron Letters

DOI:
[10.1016/S0040-4039\(01\)81740-8](https://doi.org/10.1016/S0040-4039(01)81740-8)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1984

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Vries, J. G. D., Hauser, G., & Sigmund, G. (1984). Synthesis of Carbapenems with Carbon Substituents at C-3 Using a Wittig Approach. *Tetrahedron Letters*, 25(52). [https://doi.org/10.1016/S0040-4039\(01\)81740-8](https://doi.org/10.1016/S0040-4039(01)81740-8)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SYNTHESIS OF CARBAPENEMS WITH CARBON SUBSTITUENTS


AT C-3 USING A WITTIG APPROACH

Johannes G. de Vries*, Gerhard Hauser and Gerhard Sigmund

SANDOZ Forschungsinstitut, Brunnerstraße 59, A-1235 Vienna, Austria

Summary: Reaction of ketone **2** with phosphorane **4** produced a mixture of endocyclic and exocyclic carbapenem esters **5** and **6**. Hydrogenolysis of these gave carbapenems **7** and **8** or carbapenams **9** and **10**.

Carbapenems are an important new class of antibiotics of which thienamycin (**1**) is the most well-known example¹. They distinguish themselves from the classical β -lactams such as penicillins and cephalosporins by their broad-spectrum antibacterial activity, covering a wide range of both gram-positive and gram-negative bacteria. One of the main obstacles on their way to useful therapeutic agents is their instability towards renal dehydropeptidase leading to low urinary recovery². In an attempt to identify carbapenems with greater metabolic stability we have synthesized carbapenems with fluorine instead of hydroxyl in the 8-position³. In this communication we report our synthetic efforts towards syntheses of carbapenems that in addition are devoid of sulphur in the side chain at C-3. Published syntheses of C-3 carbon substituted carbapenems⁴ have always relied on variations on the method originally developed by Woodward and coworkers⁵.



We were intrigued by the possibility of using bicyclic ketone **2** as precursor for this class of carbapenems. This compound (i.e. **2a**) was developed by Merck chemists as intermediate for the syntheses of sulphur substituted carbapenems⁶; in our laboratories we have used **2b** and **2c** for the same purpose³. The ketone moiety in **2** is exceptionally reactive for its kind: We found that most nucleophiles react instantaneously with **2b,c** giving rise to ring-opened products of type **3** (Scheme 1)⁷. We were therefore gratified to find that stabilized phosphoranes **4** reacted with **2b,c** to produce an inseparable mixture of 3-substituted carbapenem esters **5** (endo) and **6** (exo; E and Z)^{8,9} (Scheme 2, for conditions and yields see Table 1).



Table 1: Reaction of bicyclic ketone **2** with phosphoranes **4***

Entry	R	Z ¹¹	Conditions	Yield**	Product Ratio ¹⁰			Suffix
					5	6-E	6-Z	
1	CH ₃	CN	CH ₂ Cl ₂ , RT, 20 hrs.	68 %	51	36	13	k
2	CH ₃	CO ₂ Et	CH ₂ Cl ₂ , RT, 2 hrs.	94 %	83	12	5	l
3	H	CN	CH ₂ Cl ₂ , RT, 2.5 hrs.	82 %	56	34	9	m
4	H	CO ₂ Et	CH ₂ Cl ₂ , RT, 0.75 hrs.	83 %	93	5	2	n
5	H	COCH ₃	CH ₂ Cl ₂ , reflux 5 hrs.	43 %	38	62	--	o

* 1.5 eq of phosphorane, c = 0.05 - 0.07 mol/l

** Yields of purified products (column chromatography: silica, CH₂Cl₂).

With phosphoranes **4p** and **4q** the reaction proceeded very smoothly at room temperature. Reaction between **2b** and **4r** was more sluggish and took 5 hrs. at CH₂Cl₂ reflux temperature. Reactions with more basic phosphoranes such as Ph₃P=CHCl¹² and Ph₃P=CHSCH₃¹³ did not lead to products and bicyclic ketone **2** was recovered in good yield; presumably hydrogen exchange occurred.



Unfortunately, hydrogenolytic removal of the p-nitrobenzyl protective ester caused serious problems. Only with the cyanomethyl substituted carbapenem esters (5,6k and 5,6m) were we able to obtain carbapenem potassium salts 7 and 8 as inseparable mixtures in moderate yields, when short hydrogenation times were used (Table 2).

Table 2: Hydrogenolysis of carbapenem p-nitrobenzyl esters (5,6)*

Entry	R	Z	Time	Yield	Product Ratio					Suffix
					<u>7</u>	<u>8E</u>	<u>8Z</u>	<u>9</u>	<u>10</u>	
1	CH ₃	CN	1 hr.	43 %	35	53	12	traces		k
2	CH ₃	CO ₂ Et	1.25 hr.	73 %	--	--	--	21	79	l
3	H	CN	5 min.	45 %	15	55	30	traces		m
4	H	CN	1 hr.	60 %		traces		94	6	m
5	H	CO ₂ Et	15 min.	22 %		traces		65	32	n

* Conditions: 100 mg of the mixture of 5 and 6 and 100 mg of 10 % Pd/C in a mixture of 8 ml of EtOAc and 8 ml of phosphate buffer pH 7 (c = 0.15 M) were hydrogenated at 1 bar H₂ over the period indicated in the table. After filtration the aqueous layer was concentrated and purified over a C-18 column.

However, the already unfavorable endocyclic/exocyclic ratio shifted even further towards the - presumably antibacterially inactive - exocyclic product. Attempted equilibration of the product of entry 3 (Table 2) in D₂O at room temperature over 3 hrs. had the adverse effect and led to a further increase in 8m. Short hydrogenolysis of ethoxycarbonylmethyl substituted carbapenem esters (5,6l and 5,6n) only gave a mixture of carbapenamams 9 and 10¹⁴. Carbapenamams were also produced upon prolonged hydrogenation of the 3-cyanomethyl carbapenem esters 5,6m.

None of the product mixtures of Table 2 showed interesting antibacterial activity.

ACKNOWLEDGEMENT: We thank Dr. C.P. Mak (SFI) for discussions, Dr. G. Schulz (SFI) for interpretation of NMR spectra and Dr. J. Hildebrandt (SFI) for *in vitro* testing.

REFERENCES

1. R.W. Ratcliffe and G. Albers-Schönberg in "Chemistry and Biology of β -lactam Antibiotics", Vol 2, p 227, edited by R.B. Morin and M. Gorman, Academic Pres, 1982.
2. (a) H. Kropp, J.G. Sundelof, R. Hajdu and F.M. Kahan, *Antimicrob. Agents Chemother.* **1982**, 22, 62 (b) J.S. Kahan, F.M. Kahan, R. Goegelman, S.A. Currie, M. Jackson, E.O. Stapley, T.W. Miller, A.K. Miller, D. Hendlin, S. Mochales, S. Hernandez, H.B. Woodruff, and J. Birnbaum, *J. Antibiot.*, **1979**, 32, 1.
3. (a) C.P. Mak, K. Wagner, C. Mayerl, H. Fliri, *Heterocycles*, **1982**, 19, 1399. (b) C.P. Mak, H. Fliri, Belgian Patent 897351-A, 1982
4. (a) S. Oida, A. Yoshida, E. Ohki, *Chem. Pharm. Bull.* **1980**, 28, 3494. (b) J.H. Bateson, A.J.G. Baxter, P.M. Roberts, T.C. Smale, R. Southgate, *J. Chem. Soc. Perkin I*, **1981**, 3242. (c) C.L. Branch, M.J. Pearson, *J. Chem. Soc. Perkin I*, **1982**, 2123. (d) L.D. Cama, K.J. Wildonger, R. Guthikonda, R.W. Ratcliffe, B.G. Christensen, *Tetrahedron*, **1983**, 39, 2531. (e) D.H. Shih, J.A. Fayter, B.G. Christensen, *Tetrahedron Lett.*, **1984**, 1639. (f) M. Shiozaki, N. Ishida, H. Maruyama, T. Hiraoka, S. Sugawara, *J. Antibiot.* **1984**, 37, 57. (g) W. Koller, A. Linkies, H. Pietsch, H. Rehling, D. Reuschling, *Angew. Chem.*, **1982**, 94, 546. (h) E. Götschi, German Patent 3 201 957 A1, 1982.

5. R. Scartazzini, H. Peter, H. Bickel, K. Heusler, R.B. Woodward, *Helv. Chim. Acta*, **1972**, 55, 408.
6. D.G. Melillo, I. Shinkai, T. Liu, K. Ryan, M. Sletzing, *Tetrahedron Lett.*, **1980**, 2783
7. J.G. de Vries, G. Hauser, G. Sigmund, *Heterocycles*, submitted for publication.
8. All compounds are racemic mixtures
9. Selected NMR-data (only fully described for 5n, 5l, 9m and 10l. Spectra of PNB esters in CDCl_3 , salts in D_2O): 5n: δ 8.24 and 7.85 (4H, AA'BB', $J=9.0\text{Hz}$, Ar), 5.47 and 5.26 (2H, AB-q, $J=14.0\text{Hz}$, CH_2Ar), 5.00 (1H, ddq, $J=48.3$, 7.2, 6.3Hz, HCF), 4.26 (1H, ddd, $J=7.3$, 6.4, 2.6Hz, H-5), 4.17 (2H, q, $J=7.1\text{Hz}$, OCH_2), 3.82 and 3.63 (2H, AB-q, $J=16.5\text{Hz}$, $=\text{C}-\text{CH}_2$), 3.35 (1H, ddd, $J=19.0$, 7.2, 2.6Hz, H-6), 3.08 and 3.00 (2H, ABX, $J_{\text{AB}}=18.5\text{Hz}$, $J=7.3$, 6.4Hz, H-4), 1.50 (3H, dd, $J=23.8$, 6.3Hz, CH_3CF), 1.26 (3H, t, $J=7.1\text{Hz}$, CH_3CH_2). 5m: 3.94 and 3.76 (2H, AB-q, t, $J_{\text{AB}}=18.5\text{Hz}$, $J=1.5\text{Hz}$, CH_2CN). 6m-E: 5.65 (1H, dt, $J=3.1$, 2.2Hz, $=\text{CH}$), 5.16 (1H, m, H-2). 6m-Z: 5.60 (1H, dt, $J=1.6$, 2.0Hz, $=\text{CH}$), 5.40 (1H, m, H-2). 5o: 3.91 and 3.75 (2H, AB-q, $J=17.5\text{Hz}$, CH_2Ac), 2.14 (3H, s, COCH_3). 6o-E: 6.07 (1H, m, $=\text{CH}$), 4.60 (1H, m, H-2). 5l: 8.23 and 7.66 (4H, AA'BB', $J=9.0\text{Hz}$, Ar), 5.49 and 5.26 (2H, AB-q, $J=14.5\text{Hz}$, CH_2Ar), 4.29 (1H, ddd, $J=9.8$, 8.5, 3.3Hz, H-5), 4.16 (2H, q, $J=7.2\text{Hz}$, OCH_2), 3.92 and 3.84 (2H, AB-q, $J=16.5\text{Hz}$, $=\text{C}-\text{CH}_2$), 3.36 (1H, dd, $J=22.5$, 3.3Hz, H-6), 3.09 and 2.97 (2H, ABX, $J_{\text{AB}}=19.0\text{Hz}$, $J=9.8$, 8.5Hz, H-4), 1.57 (3H, d, $J=21.3\text{Hz}$, CH_3CF), 1.48 (3H, d, $J=21.3\text{Hz}$, CH_3CF), 1.26 (3H, t, $J=7.2\text{Hz}$, CH_2CH_3). 5k: 3.95 and 3.79 (2H, AB-q, t, $J_{\text{AB}}=18.5\text{Hz}$, $J=1.5\text{Hz}$, CH_2CN). 6k-E: 5.65 (1H, dt, $J=3.0$, 2.0Hz, $=\text{CH}$), 5.19 (1H, m, H-2), 6k-Z: 5.59. (1H, dt, $J=1.5$, 2.5Hz, $=\text{CH}$), 5.41 (1H, m, H-2). 7m: 3.94 and 3.81 (2H, AB-q, $J=18.5\text{Hz}$, CH_2CN). 8m-E: 5.81 (1H, dt, $J=3.0$, 2.0Hz, $=\text{CH}$), 5.02 (1H, m, H-2), 8m-Z: 5.71 (1H, dt, $J=2.0$, 2.2Hz), 5.16 (1H, m, H-2). 7k: 4.29 (1H, ddd, $J=9.5$, 8.5, 3.0Hz, H-5), 3.95 and 3.80 (2H, AB-q, t, $J_{\text{AB}}=18.3\text{Hz}$, $J=1.2\text{Hz}$, CH_2CN), 3.68 (1H, dd, $J=27.1$, 3.0Hz, H-6). 8k-E: 5.81 (1H, dt, $J=3.0$, 2.0Hz, $=\text{CH}$), 5.02 (1H, m, H-2), 4.19 (1H, ddd, $J=7.0$, 7.0, 2.1Hz, H-5), 3.53 (1H, dd, $J=27.0$, 2.1Hz, H-6). 8k-Z: 5.71 (1H, dt, $J=2.0$, 2.2Hz, $=\text{CH}$), 5.16 (1H, m, H-2), 3.52 (1H, dd, $J=26.6$, 2.3Hz, H-6). 9n: 4.45 (1H, d, $J=7.4\text{Hz}$, H-2), 4.04-4.21 (5H, m, CH_2CO , H-5), 3.38 (1H, ddd, $J=27.3$, 4.5, 2.5Hz, H-6), 3.02-3.21 (1H, m, H-3). 10n: 3.92 (1H, d, $J=8.5\text{Hz}$, H-2), 9m: 5.15 (1H, ddq, $J=47.8$, 5.0, 6.3Hz, HCF), 4.47 (1H, d, $J=7.0\text{Hz}$, H-2), 4.16 (1H, ddd, $J=6.3$, 6.3, 2.5Hz, H-5), 3.41 (1H, ddd, $J=27.3$, 5.0, 2.5Hz, H-6), 3.14 (1H, m, H-3), 2.65 and 2.60 (2H, ABX, $J_{\text{AB}}=17.5\text{Hz}$, $J=9.3$, 6.0Hz, CH_2CN), 2.26 and 2.15 (2H, ABXY, $J_{\text{AB}}=15.5\text{Hz}$, $J_{2,4}=J_{2,4'}=J_{3,4}=J_{3,4'}=6.3\text{Hz}$, H-4), 1.45 (3H, dd, $J=24.8$, 6.3Hz, CH_3CF). 9l: 4.44 (1H, d, $J=7.3\text{Hz}$, H-2), 3.40 (1H, dd, $J=27.0$, 2.8Hz, H-6). 10l: 4.18 (2H, q, $J=7.1\text{Hz}$, OCH_2), 3.89 (1H, d, $J=7.8\text{Hz}$, H-2), 3.87 (1H, ddd, $J=9.8$, 5.3, 2.3Hz, H-5), 3.42 (1H, dd, $J=27.3$, 2.3Hz, H-6), 3.15 (1H, m, H-3), 2.59 and 2.52 (2H, ABX, $J_{\text{AB}}=16.3\text{Hz}$, $J=7.8$, 7.5Hz, CH_2CO_2), 2.14-2.24 (1H, m, H-4), 1.62-1.76 (1H, m, H-4), 1.53 (3H, d, $J=22.5\text{Hz}$, CH_3CF), 1.45 (3H, d, $J=22.5\text{Hz}$, CH_3CF), 1.27 (3H, t, $J=7.1\text{Hz}$, CH_3CH_2). All other spectral data were in agreement with the proposed structures.
10. These were not the thermodynamic equilibria ratios. In most occasions the ratios changed over longer periods of time. Determined by NMR and HPLC.
11. Prepared from the corresponding phosphonium salt and aq. K_2CO_3 .
12. R. Appel and W. Morbach, *Angew. Chem.* **1977**, 89, 203.
13. R. Greenwald, M. Chaykovsky, E.J. Corey, *J. Org. Chem.* **1963**, 28, 1128
14. Stereochemical assignment by comparison of NMR data with those published for carbapenamams of known stereochemistry: J.H. Bateson, P.M. Roberts, T.C. Smale and R. Southgate, *J. Chem. Soc. Chem. Commun.* **1980**, 185; T. Haneishi, M. Nakajima, N. Serizawa, M. Inukai, Y. Takiguchi, M. Arai, S. Satoh, H. Kuwano, C. Tamura, *J. Antibiot.* **1983**, 36, 1581.

(Received in Germany 18 September 1984)